

Diagnostic Value of Thyrotropin-Releasing-Hormone Stimulation in Patients With Pituitary Tumor

DAVID M. COOK, MD; MONTE A. GREER, MD, and H. PAXTON, MD, Portland

Plasma prolactin response to thyrotropin-releasing-hormone (TRH) stimulation was diminished in 30 patients with prolactinomas and 9 patients with acromegaly who had normal serum prolactin levels. There was no overlap of prolactin responses when compared with 32 control patients. Responses of ten patients with adrenocorticotropin (ACTH)-secreting pituitary tumors were similar to those of controls. Plasma growth hormone concentrations after TRH stimulation changed significantly in 28% of normal control and 20%, 25% and 50% of patients with prolactin-, growth hormone- and ACTH-secreting pituitary tumors, respectively. Our data suggest that the blunted TRH-induced rise in plasma prolactin levels in patients with prolactinomas and those with acromegaly may be related to humoral factor(s) affecting TRH receptor or postreceptor function. Growth hormone responses to TRH are nonspecific and should not be considered a marker for active acromegaly.

(Cook DM, Greer MA, Paxton H: Diagnostic value of thyrotropin-releasing-hormone stimulation in patients with pituitary tumor. West J Med 1987 Aug; 147:161-164)

The exact mechanism(s) responsible for the blunted plasma prolactin responses to thyrotropin-releasing-hormone (TRH) stimulation in patients with prolactinomas is unknown. TRH receptors have been found on human prolactinoma cells.¹ In vitro responses of these cells to TRH administration have been variable.^{2,3} In most patients with hyperprolactinemia not caused by a pituitary tumor, there is a normal prolactin secretory response to TRH,⁴⁻⁸ suggesting that ultrashort-loop feedback of prolactin on lactotrope activity is not responsible for the decreased response in prolactinoma patients.

We wondered if the presence of any hypersecreting pituitary tumor might blunt responses to TRH. A tumor mass might impair the local blood supply and thus the delivery of TRH to prolactin-secreting cells, accounting for inadequate responses. Patients with prolactinomas were compared with patients with adrenocorticotropin (ACTH)- and growth hormone-secreting tumors. We found the responses of patients with acromegaly were similar to those of patients with prolactinomas, and patients with ACTH-secreting tumors had responses similar to those of controls. Our studies suggest humoral and not mass lesion effects may explain blunted responses and that growth hormone- and prolactin-secreting tumors are capable of inhibiting prolactin responses to TRH, possibly by a similar mechanism.

Patients and Methods

Controls

Six men and 26 women were studied initially because abnormal pituitary function was questioned clinically. Typical problems included mild obesity (possible Cushing's disease), coarse facial features (possible acromegaly) and possibly a partially eroded sella (verified later as normal by

computed tomographic [CT] scan). All findings later proved to be normal except in two patients with primary hypothyroidism and one with drug-induced (thioridazine hydrochloride [Mellaril]) hyperprolactinemia. The two patients with hypothyroidism had pituitary enlargement and hyperprolactinemia: pituitary size and prolactin concentrations reverted to normal with thyroxine therapy.

Prolactinoma

In all, 23 women and 7 men had prolactinomas verified by a surgical procedure. All patients had elevated basal plasma prolactin concentrations. Plasma growth hormone concentrations were normal and suppressed with glucose. All were studied during the preoperative period.

Acromegaly

Four men and five women had acromegaly associated with elevated growth hormone concentrations not suppressible by glucose. None had hyperprolactinemia. All were studied preoperatively.

Cushing's Disease

Ten patients with ACTH-secreting pituitary tumors had detectable levels of plasma ACTH. The diagnosis was established by standard dexamethasone suppression and metyrapone stimulation tests. Plasma growth hormone concentrations were suppressed by glucose in all patients to undetectable levels. All tumors were verified by a pituitary operation. All patients except one were cured by surgical therapy.

All patients were admitted to the Clinical Research Center of the University of Oregon Health Sciences Center. Testing was done between 8 AM and 9 AM in the fasting basal state. After informed consent, 500 µg TRH was given as an intrave-

ABBREVIATIONS USED IN TEXT

ACTH = adrenocorticotropin

CT = computed tomography

RIA = radioimmunoassay

TRH = thyrotropin-releasing hormone

nous bolus via an indwelling venous catheter. A blood specimen was taken at 0, 20 and 40 minutes through the same catheter and assayed for prolactin and growth hormone by described techniques. Growth hormone was assayed by double-antibody radioimmunoassay (RIA) (BIO-RIA Incorporated, Montreal, Canada). Prolactin was assayed by double-antibody RIA (Diagnostic Products Corporation, Los Angeles).

Results

Prolactin responses to 500 μ g TRH are shown in Figure 1. Peak responses were predominantly at 20 minutes, but those occurring at 40 minutes are not identified separately.

Peak plasma prolactin responses are expressed as a percentage increase from basal. TRH-induced changes in plasma prolactin concentrations were significantly different when patients with prolactinomas were compared with either controls ($P < .00001$) or patients with ACTH-secreting tumors




Figure 1.—The percent change in plasma prolactin levels in response to the administration of 500 μ g of thyrotropin-releasing hormone preoperatively is shown. Basal prolactin concentrations in the prolactinoma group ranged from 22 to 1,560 ng per ml. Two patients in the control group with primary hypothyroidism are identified by triangles and have basal prolactin concentrations of 10 and 34 ng per ml, respectively. The basal prolactin concentration of the one patient with drug-induced hyperprolactinemia (identified by a square) was 95 ng per ml. Basal prolactin concentrations were less than 20 ng per ml in all patients with growth hormone (GH)- and adrenocorticotropin (ACTH)-secreting tumors and in the remaining control patients.

($P < .00001$; Kruskal-Wallis analysis of variance). Patients with growth hormone-secreting tumors had significantly different responses from those of controls ($P < .00001$). There was no significant difference in responses between patients with prolactinoma and those with acromegaly ($P < .5$). Responses expressed in nanograms per milliliter are shown in Figure 2 for all patients studied.

Individual basal and peak plasma growth hormone concentrations after TRH stimulation in the four groups are shown in Figure 3. A growth hormone response is defined arbitrarily as a level rising 100% or more above basal and a 5-ng-per-ml rise because spontaneous fluctuations could account for large percentage changes if basal values are low. Significant growth hormone responses to TRH stimulation in each group were as follows: control group 28%, prolactinoma group 20%, acromegalic group 25% and group with ACTH-secreting tumors 50%.

Postoperative Response

Not all patients were cured (Figure 4). The criteria for cure in patients with prolactinomas included normal prolactin




Figure 2.—The graphs show peak plasma prolactin responses (ng per ml) to thyrotropin-releasing hormone stimulation. The two patients in the control group with primary hypothyroidism are identified by triangles. The one patient with drug-induced hyperprolactinemia is identified by a square. ACTH = adrenocorticotropin, B = basal, GH = growth hormone, P = peak




Figure 3.—Plasma growth hormone responses to the administration of 500 μ g of thyrotropin-releasing hormone in a preoperative study are given. Absolute values are shown and suggest significant responses in all four groups. ACTH = adrenocorticotropin, B = basal, GH = growth hormone, P = peak

levels on at least three separate determinations and no evidence of an anterior pituitary deficiency. Patients not cured of prolactinoma included those in whom the surgeon believed not all the tumor was removed and patients who had persistent hyperprolactinemia. None of the noncured patients were thought to have hyperprolactinemia on the basis of isolation of the anterior pituitary from the hypothalamus. Prolactin responses to TRH in patients with prolactinoma and those with acromegaly were similar to those of controls in cured patients. Those not cured were persistently unresponsive. Patients with ACTH-secreting tumors were unchanged postoperatively whether cured or not. Not all patients returned for follow-up studies.

Discussion

Abnormal prolactin responses to TRH have been observed in most patients with prolactinomas.* Klijn and co-workers have suggested that this may be related to a tumor in the hypothalamic area because 7 of 11 patients they studied with suprasellar tumors that were not prolactinomas had blunted responses.²⁰ Our data suggest that other mechanism(s) for blunted responses may exist in growth hormone- and prolactin-secreting tumors.

It has been suggested that patients with prolactinomas may have a benign course.²¹⁻²³ Such studies that suggest spontaneous regression or stable growth characteristics, however, lack adequate proof that these patients actually have tumors. A prolactinoma may or may not be visible by high-resolution CT scan of the sella.²⁴⁻²⁸ A reliable nonoperative means of identifying tumors would avoid an unnecessary operation and the undue worry if a tumor was not present and improve our

knowledge of the natural history of this tumor. Our results suggest that prolactinoma patients do not respond to TRH stimulation with a change in plasma prolactin concentrations. Most series reporting false-positive responses do not have surgical and pathologic verification of prolactin-secreting tumors, and those that do have histologic proof tend to implicate TRH hyporesponsiveness.⁷⁻¹⁹ Chlorpromazine stimulation tests have also been used to separate patients with prolactinomas from those with functional hyperprolactinemia. Boyd and associates found blunted prolactin responses to chlorpromazine stimulation in 4 of 6 prolactinoma patients and in 4 of 12 other patients with galactorrhea without evidence of adenoma.¹⁵ Kleinberg and colleagues found blunted rises of serum prolactin levels after chlorpromazine stimulation in all nine patients with verified prolactinoma and in 19 of 36 patients with other forms of galactorrhea.²⁹ Zárate and co-workers showed normal responses to chlorpromazine in 8 of 16 patients with galactorrhea and amenorrhea, including 2 of 5 with pituitary tumors.³⁰ Our studies with TRH would suggest a lower incidence of false-positive responses compared with those with chlorpromazine.

Why patients with prolactinomas respond poorly to TRH stimulation is an unanswered question. Patients with hyperprolactinemia of a clearly secondary cause such as drug-induced or associated with hypothyroidism usually respond to TRH administration.^{15,29,31} Other patients identified as functional may or may not respond.^{7,11,12} Because this functional group has not had surgical exploration, we cannot be sure whether or not failure to respond may represent the presence of a tumor. Our series includes two patients with hyperprolactinemia due to hypothyroidism and one due to drug-related causes. All these responded vigorously to TRH stimulation. Our series of prolactinoma patients is important because all were clearly verified by a surgical procedure.

The finding of a persistent blunted response postoperatively in only the patients with prolactinoma and acromegalic patients not cured by surgical therapy suggests that the blunted response is not of hypothalamic origin or a generalized abnormality of lactotropes. It does suggest that a complete cure is possible by surgical treatment and that a return of prolactin responsiveness to TRH stimulation may be a marker for a surgical cure.

The diminished prolactin response to TRH administration in patients with prolactinoma and acromegalic patients is not explained at present. The absence of stored prolactin in tumor could be an explanation. We did not have immunocytochemistry staining to verify the presence or absence of prolactin in tumor cells. Clear responses in the secondary causes, however, suggest that some hyperprolactinemic states are not associated with exhaustion of stored prolactin. Pathologic studies using immunocytochemistry have verified the presence of stored hormone in some, but not all, prolactinomas. We do not know of a study showing a correlation between the absence of storage and blunted or absent prolactin responses to TRH. Until such studies are forthcoming, we will continue to look for the presence of a prolactinoma when serum prolactin responses to TRH stimulation are low or absent.

Hyperprolactinemic patients with chronic renal failure also have poor plasma prolactin responses to TRH.³²⁻³⁵ Normal responsiveness does not return with dialysis but does with transplantation,³⁶ suggesting an inhibition at the pituitary level by a humoral factor other than prolactin because other functional causes of excess prolactin secretion such as

*References 7-13 (p 225), 14-19 (p 131).




Figure 4.—Plasma responses to thyrotropin-releasing hormone stimulation reverted to normal in cured patients with prolactinomas and those with acromegaly. The responses in noncured patients in both groups were unchanged. ACTH = adrenocorticotropin, C = cured, GH = growth hormone, NC = not cured

drug-induced and primary hypothyroidism are not associated with inhibition.

The growth hormone responses to TRH observed in both control patients and those with a pituitary tumor without acromegaly were surprising. Although previous studies^{35,37,38} of growth hormone responses to TRH in controls disagree with our findings, closer inspection of these reports (especially that of Irie and Tsushima³⁷) suggests that growth hormone may be significantly released by TRH stimulation in control subjects. Our findings imply that growth hormone response to TRH is not an exclusive tumor marker for acromegaly as 28% of the control patients, 20% of the patients with prolactinomas and 50% of patients with ACTH-secreting tumors had significant increases in growth hormone levels. Therefore, we suggest that the current clinical practice of accepting growth hormone responses to TRH stimulation^{37,39-42} as evidence for acromegaly may not be valid.

REFERENCES

1. Le Dafniet M, Brandi AM, Bression D, et al: Evidence of receptors for thyrotropin-releasing hormone in human prolactin-secreting adenomas. *J Clin Endocrinol Metab* 1983; 57:425-427
2. Chihara K, Iwasaki J, Minamitani N, et al: Prolactin secretion from human prolactinomas perfused in vitro: Effect of TRH, prostaglandin E₁, theophylline, dopamine and dopamine receptor blockers. *Acta Endocrinol (Copenh)* 1984; 105:6-13
3. Adams EF, Brajkovic IE, Mashiter K: Hormone secretion by dispersed cell cultures of human pituitary adenomas: Effects of theophylline, thyrotropin-releasing-hormone, somatostatin, and 2-bromo-ergocryptine. *J Clin Endocrinol Metab* 1979; 49:120-126
4. Ferrari C, Rampini P, Benco R, et al: Functional characterization of hypothalamic hyperprolactinemia. *J Clin Endocrinol Metab* 1982; 55:897-901
5. Caro JF, Israel HL, Glennon JA: Galactorrhea in sarcoidosis: Dynamic studies of prolactin, growth and gonadotropic hormone levels. *Am J Med Sci* 1979; 277:289-294
6. Marrs RP, Bertolli SJ, Kletzky OA: The use of thyrotropin-releasing hormone in distinguishing prolactin-secreting pituitary adenoma. *Am J Obstet Gynecol* 1980; 138:620-625
7. Cowden EA, Ratcliffe JG, Thomson JA, et al: Tests of prolactin secretion in diagnosis of prolactinomas. *Lancet* 1979; 1:1155-1158
8. Assies J, Schellekens APM, Touber JL: The value of an intravenous TRH test for the diagnosis of tumoural hyperprolactinaemia. *Acta Endocrinol (Copenh)* 1980; 94:439-449
9. Lamberts SWJ, Birkenhäger JC, Kwa HG: Basal and TRH-stimulated prolactin in patients with pituitary tumours. *Clin Endocrinol (Oxf)* 1976; 5:709-711
10. Ayalon D, Persitz E, Ravid R, et al: The diagnostic value of pharmacodynamic tests in the hyperprolactinaemic syndrome. *Clin Endocrinol (Oxf)* 1979; 11:201-215
11. Schaison G, Nathan C, L'Hermite M, et al: Etude de la prolactine et des gonadotrophines plasmatiques dans les aménorrhées-galactorrhées—Exploration dynamique dans 13 cas. *Nouv Presse Med* 1975; 4:644-648
12. Lachelin GCL, Abu-Fadil S, Yen SSC: Functional delineation of hyperprolactinemic amenorrhea. *J Clin Endocrinol Metab* 1977; 44:1163-1174
13. Faglia G, Beck-Peccoz P, Travaglini P, et al: Functional studies in hyperprolactinemic states. In Crosignani PG, Robyn C (Eds): *Prolactin and Human Reproduction*. London, Academic Press, 1977
14. Barbarino A, de Marinis L, Menini E, et al: Prolactin-secreting pituitary adenomas: Prolactin dynamics before and after transphenoidal surgery. *Acta Endocrinol (Copenh)* 1979; 91:397-409
15. Boyd AE III, Reichlin S, Turksoy NR: Galactorrhea-amenorrhea syndrome: Diagnosis and therapy. *Ann Intern Med* 1977; 87:165-175
16. Schwinn G, von zur Mühlen A, Köbberling J, et al: Plasma prolactin levels after TRH and chlorpromazine in normal subjects and patients with impaired pituitary function. *Acta Endocrinol (Kbh)* 1975; 79:663-676
17. Samaan NA, Leavens ME, Jesse JH Jr: Serum prolactin in patients with 'functionless' chromophobe adenomas before and after therapy. *Acta Endocrinol (Kbh)* 1977; 84:449-460
18. Jaquet P, Grisoli F, Guibout M, et al: Prolactin secreting tumors—Endocrine status before and after surgery in 33 women. *J Clin Endocrinol Metab* 1978; 46:459-466
19. Silvestrini F, Liuzzi A, Chiodini PG: Prolactin and pituitary tumors. In Martini L, James VHT (Eds): *Current Topics in Experimental Endocrinology*. New York and London, Academic Press, 1978
20. Klijn JGM, Lamberts SWJ, de Jong FH, et al: The value of thyrotropin-releasing hormone test in patients with prolactin-secreting pituitary tumors and suprasellar non-pituitary tumors. *Fertil Steril* 1981; 35:155-161
21. Koppelman MCS, Jaffe MJ, Rieth KG, et al: Hyperprolactinemia, amenorrhea, and galactorrhea—A retrospective assessment of twenty-five cases. *Ann Intern Med* 1984; 100:115-121
22. March CM, Kletzky OA, Davajan V, et al: Longitudinal evaluation of patients with untreated prolactin-secreting pituitary adenomas. *Am J Obstet Gynecol* 1981; 139:835-844
23. Cassar J, Yeung RTT, Gonticas SC, et al: Long-term follow-up of an untreated patient with probable prolactinoma (Letter). *N Engl J Med* 1984; 311:332
24. Banna M, Baker HL Jr, Houser OW: Pituitary and parapituitary tumours on computed tomography—A review article based on 230 cases. *Br J Radiol* 1980; 53:1123-1143
25. Gardeur D, Naidich TP, Metzger J: CT analysis of intrasellar pituitary adenomas with emphasis on patterns of contrast enhancement. *Neuroradiology* 1981; 20:241-247
26. Daniels DL, Williams AL, Thornton RS, et al: Differential diagnosis of intrasellar tumors by computed tomography. *Radiology* 1981; 141:697-701
27. Taylor S: High resolution computed tomography of the sella. *Radiol Clin North Am* 1982; 20:207-235
28. Wolpert SM, Post KD, Biller BJ, et al: The value of computed tomography in evaluating patients with prolactinomas. *Radiology* 1979; 131:117-119
29. Kleinberg DL, Noel GL, Frantz AG: Galactorrhea: A study of 235 cases, including 48 with pituitary tumors. *N Engl J Med* 1977; 296:589-600
30. Zárate A, Jacobs LS, Canales ES, et al: Functional evaluation of pituitary reserve in patients with the amenorrhea-galactorrhea syndrome utilizing luteinizing hormone-releasing hormone (LH-RH), L-dopa and chlorpromazine. *J Clin Endocrinol Metab* 1973; 37:855-859
31. Aono T, Shioji T, Kinugasa T, et al: Clinical and endocrinological analysis of patients with galactorrhea and menstrual disorders due to sulpiride or metaclopramide. *J Clin Endocrinol Metab* 1978; 47:675
32. Sievertsen GD, Linn VS, Nakawatase C, et al: Metabolic clearance and secretion rates of human prolactin in normal subjects and in patients with chronic renal failure. *J Clin Endocrinol Metab* 1980; 50:846-852
33. Gomez F, Cueva R, Wauters JP, et al: Endocrine abnormalities in patients undergoing long term hemodialysis—The role of prolactin. *Am J Med* 1980; 68:522-530
34. Weetman AP, Weightman DR, Scanlon MF: Impaired dopaminergic control of thyroid stimulating hormone secretion in chronic renal failure. *Clin Endocrinol (Oxf)* 1981; 15:451-456
35. Gonzalez-Barcena D, Kastin AJ, Schalch DS, et al: Responses to thyrotropin-releasing hormone in patients with renal failure and after infusion in normal men. *J Clin Endocrinol Metab* 1973; 36:118-120
36. Lim JS, Kathpalia SC, Frohman LA: Hyperprolactinemia and impaired pituitary response to suppression and stimulation in chronic renal failure: Reversal after transplantation. *J Clin Endocrinol Metab* 1979; 48:101-106
37. Irie M, Tsushima T: Increase of serum growth hormone concentration following thyrotropin-releasing hormone injection in patients with acromegaly or gigantism. *J Clin Endocrinol Metab* 1972; 35:97-100
38. Maeda K, Kato Y, Ohgo S, et al: Growth hormone and prolactin release after injection of thyrotropin-releasing hormone in patients with depression. *J Clin Endocrinol Metab* 1975; 40:501-505
39. Liuzzi A, Chiodini PG, Botalla L, et al: Growth hormone (GH)-releasing activity of TRH and GH-lowering effect of dopaminergic drugs in acromegaly: Homogeneity in the two responses. *J Clin Endocrinol Metab* 1974; 39:871-876
40. Faglia G, Paracchi A, Beck-Peccoz P, et al: An explanatory hypothesis for plasma GH response to non-specific releasing hormones and for paradoxical GH inhibition after dopaminergic drugs in acromegaly. *Acta Endocrinol (Copenh)* [Suppl 199] 1975; 80:323
41. Winkelmann W: Plasma growth hormone response to TRH, LHRH and bromocriptine in patients with acromegaly. *Acta Endocrinol (Copenh)* [Suppl 208] 1977; 84:26-27
42. Lamberts SWJ, Klijn JGM, Kwa GH, et al: The dynamics of growth hormone and prolactin secretion in acromegalic patients with 'mixed' pituitary tumours. *Acta Endocrinol (Copenh)* 1979; 90:198-210